



- N.B.: (1) Question No. 1 is compulsory  
(2) Attempt any Three from remaining

- Q1 a) If  $X_1$  has mean 4 and variance 9 &  $X_2$  has mean  $-2$  and variance 4 [5]  
where  $X_1$  &  $X_2$  are independent, find  $E(2X_1 + X_2 - 3)$  and  
 $V(2X_1 + X_2 - 3)$ .
- b) Find the extremals of  $\int_{x_1}^{x_2} (x + y')y' dx$  [5]
- c) Verify Cauchy Schwartz inequality for the vectors  $u = (-4, 2, 1)$  and [5]  
 $v = (8, -4, -2)$
- d) Check whether  $A = \begin{bmatrix} 2 & -2 & 3 \\ 1 & 1 & 1 \\ 1 & 3 & -1 \end{bmatrix}$  is derogatory or not. [5]
- Q2 a) Using Cauchy's Residue theorem evaluate  $\int_C \frac{z-1}{(z+1)^2(z-2)}$  where  $C$  is [6]  
 $|z| = 4$
- b) Show that the extremal of the isoperimetric problem [6]  
 $I[y(x)] = \int_{x_1}^{x_2} (y')^2 dx$  subject to the condition  $\int_{x_1}^{x_2} y dx = k$  is a  
parabola.
- c) Is the matrix  $A = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 0 & 0 & 1 \end{bmatrix}$  diagonalisable? If so find the diagonal [8]  
matrix and the transforming matrix.
- Q3 a) Verify Cayley-Hamilton theorem for  $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \\ 3 & 1 & -1 \end{bmatrix}$  [6]  
hence find  $A^{-1}$
- b) Check whether the following are subspaces of  $\mathbb{R}^3$  [6]  
(i)  $W = \{(a, 0, 0) \mid a \in \mathbb{R}\}$   
(ii)  $W = \{(x, y, z) \mid x = 1, z = 1, y \in \mathbb{R}\}$
- c) Expand  $f(z) = \frac{1}{(z-1)(z-2)}$  in Taylor's & Laurent's series indicating [8]  
regions of convergence.

Q4 a) Using Rayleigh-Ritz method to solve the boundary value problem [6]

$$I = \int_0^1 (2xy + y^2 - (y')^2) dx ; 0 \leq x \leq 1 \text{ given } y(0) = y(1) = 0$$

b) If  $A = \begin{bmatrix} -1 & 4 \\ 2 & 1 \end{bmatrix}$  then prove that  $3 \tan A = A \tan 3$ . [6]

c) If sizes of 10,000 items are normally distributed with mean 20 cms & standard deviation of 4 cms. Find the probability that an item selected at random will have size :

(i) between 18 cms and 23 cms , (ii) above 26 cms

Q5 a) Find orthonormal basis of  $\mathbb{R}^3$  using Gram-Schmidt process where [6]

$$S = \{(1,0,0) , (3,7, -2) , (0,4,1)\}$$

b) In a factory, machines A , B & C produce 30%, 50% & 20% of the total production of an item. Out of their production 80% , 50% & 10% are defective respectively. An item is chosen at random and found to be defective. What is the probability that it was produced by machine A. [6]

c) Evaluate  $\int_{-\infty}^{\infty} \frac{dx}{(x^2+4)(x^2+9)}$  [8]

Q6 a) Evaluate  $\int_C \frac{dz}{z^3(z+4)}$  where C is the circle [6]

(i)  $|z| = 2$  and (ii)  $|z - 3| = 2$

b) Two unbiased dice are thrown three times, using Binomial distribution find the probability that the sum nine would be obtained (i) once , (ii) twice [6]

c) For the following data [8]

X	100	110	120	130	140	150	160	170	180	190	
Y	45	51	54	61	66	70	74	78	85	89	

Find the coefficients of regression  $b_{xy}$  &  $b_{yx}$  and the coefficient of correlation ( $r$ )

\*\*\*\*\*